

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/340654097>

Modeling behaviour of wet and dry days in Iran from the perspective of Markov chains

Article in *Mausam* · January 2020

DOI: 10.54302/mausam.v7i1i.8

CITATION

1

READS

67

3 authors, including:



Peyman Mahmoudi

University of Sistan and Baluchestan

65 PUBLICATIONS 121 CITATIONS

[SEE PROFILE](#)



Seyed Mahdi Amir Jahanshahi

University of Sistan and Baluchestan

38 PUBLICATIONS 251 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



انتخاب بهترین توابع توزیع احتمالاتی برای محاسبه شاخص بارش استاندارد شده در مقیاس های زمانی مختلف برای ایران [View project](#)



Goodness-of-Fit Tests for Complete (Uncensored) Data [View project](#)

Modeling behaviour of wet and dry days in Iran from the perspective of Markov chains

PEYMAN MAHMOUDI, SEYED MAHDI AMIR JAHANSHAH and ZOHREH MORADI

University of Sistan and Baluchestan, Zahedan, Iran

Payame Noor University of Saghez, Saghez, Kurdistan Province, Iran

(Received 20 December 2017, Accepted 19 November 2019)

e mail : p_mahmoudi@gep.usb.ac.ir

सार – वर्तमान अध्ययन का उद्देश्य ईरान में मार्कोव चेन मॉडल का उपयोग करके आर्द्र और बिना बारिश वाले दिनों की स्थिति का मॉडल तैयार करना है। इसके लिए, ईरान मौसम विज्ञान संगठन से 25 साल के अंतराल (1991-2015) के लिए 44 सिनॉप्टिक स्टेशनों की दैनिक वर्षा से संबंधित डेटा प्राप्त किया गया। फिर, बिना बारिश वाले दिनों और आर्द्र दिनों की घटना की स्थिर संभावना सहित ईरान के बिना बारिश वाले दिनों और आर्द्र दिनों की मार्कोव विशेषताएँ, बिना बारिश वाले दिनों की अवधि की अपेक्षित लंबाई, आर्द्र अवधि की अपेक्षित लंबाई, बिना बारिश वाले दिनों-आर्द्र वाले दिनों का दौर चक्र, बिना बारिश वाले दिनों या आर्द्र एपिसोड के लिए वापसी की अवधि और अंत में 5, 10, 15, 20, 25, और 30 दिनों के लिए बिना बारिश वाले दिनों की निरंतरता संभावना की गणना एक मौसमी पैमाने में सभी सिनॉप्टिक स्टेशनों के लिए की गई। प्राप्त हुए परिणामों से पता चला कि पूरे ईरान में अलग-अलग संभावनाओं के साथ शरद ऋतु, शीत ऋतु और वसंत ऋतु के तीन मौसमों में कम शुष्कता बने रहने (5 और 10 दिन) की घटना होती है। हालांकि, मौसम और जगह के लिहाज से दीर्घकालीन बिना बारिश वाले दिनों की निरंतरताओं (20 दिनों से अधिक) की घटना की संभावना रहती है, सर्दियों में ईरान के उत्तरी आधे हिस्से में इस तरह की निरंतरताओं की घटना की कोई संभावना नहीं होती है। जैसा कि शरद ऋतु और वसंत में ईरान के ऊपरी वायुमंडल के स्तरों में वायुमंडल के दीर्घकालिक स्थिरता की स्थिति का अंत और शुरुआत होती है, विशेष रूप से ईरान के दक्षिणी आधे हिस्से में 30-दिन के बिना बारिश वाले दिनों की आवधिक घटना की संभावना बढ़ जाती है। इसके अलावा, ईरान के हर हिस्से के लिए बिना बारिश वाले दिनों के लिए अपेक्षित वापसी की अवधि लगभग स्थिर रहती है और यह 1 और 2 दिनों के बीच की सीमा में रहती है। हालांकि वर्षा अवधि के लिए वापसी दिनों की संख्या इस नियम का पालन नहीं करती है और ईरान के हर हिस्से के लिए भिन्न होती है ताकि शरद ऋतु में 2.15 दिनों से लेकर वसंत ऋतु में 79 दिनों तक बदलता रहे, जो ईरान की जलवायविक विविधता की ओर इशारा करता है।

ABSTRACT. The current study aims to model the behaviour of wet and dry days in Iran using Markov Chain Models. To this end, data related to daily precipitation of 44 synoptic stations for a 25-years interval (1991-2015) was obtained from Iran Meteorological Organization. Then, the Markov features of dry and wet days of Iran including stationary probabilities of dry and wet days occurrence, the expected length of dry periods, the expected length of wet periods, dry-wet spells cycle, return periods for dry or wet episodes and finally, the possibility of occurrence of the continuity of dry days for 5, 10, 15, 20, 25 and 30 days were calculated for all the synoptic stations in a seasonal scale. The results showed that there is the occurrence of dry short continuities (5 and 10 days) in three seasons of autumn, winter and spring with different possibilities all over Iran. However, the possibility of occurrence of long-term dry continuities (more than 20 days) is variable in terms of season and place so that in winter, no possibility of occurrence of this type of continuities is obvious in the northern half of Iran. As in autumn and spring those are the end and beginning of long-term stability conditions of the atmosphere in the upper atmosphere levels of Iran, the possibility of periodical occurrence of 30-days dry days, particularly in the southern half of Iran increases. In addition, the expected return periods for dry days is almost steady for every part of Iran and is in the range between 1 and 2 days. However, the number of return days to a precipitation period does not follow this rule and varies for every part of Iran so that from 2.15 days in autumn to 79 days in spring is variable, pointing to the climate diversity of Iran.

Key words – Dry day, Wet day, Modeling, Markov chain, Iran.

1. Introduction

Precipitation, as the most important climatic element, has had always special complexities in Iran. These complexities that are mostly due to the geographical status

of this broad land have made precipitation not to have a uniform spatial and temporal distribution. In the last few years, the precipitation features of Iran have been studied from different aspects. One of these aspects is to study the structure of wet days in Iran and its regionalization by

using multivariate statistical methods (Nazaripour, 2011). Another important aspect considered by Babaie and Farajzadeh (2002) was to study the spatial-temporal changes patterns of precipitation in Iran. Movahedi *et al.* (2012) also considered the variability of precipitation regimes by using harmonic analysis. Also, the ratio of maximum daily precipitation to Iran's annual precipitation is another subject considered by Zolfaghari *et al.* (2009).

From the perspective of water resource management, only relying on short-term and medium-term predictions of precipitation in a region cannot be enough, but a comprehensive understanding of the spatial-temporal behaviour of wet and dry periods is also considered as a necessity (Lana and Burgueno, 1998). In order to achieve this comprehensive understanding, a set of statistical methods is needed to model some important aspects like the length of consecutive dry periods or probabilities of consecutive dry days.

Since the length of dry periods is considered a discrete variable by using daily precipitation data, thus discrete models like geometric distribution or truncated negative binomial (De Arruda and Pinto, 1980) can be used to study it. However, using continuous distributions like the exponential distribution in these studies is not very unusual (for example see the study of Burgueno *et al.* (1994)). Using Markov chains, due to their abilities in comparison to other models, can give us a complete description of the behaviour of wet and dry days dependent on a meteorological station.

Among the classic references that explained this statistical method in a completed way, the studies of Kemeny & Snell (1960), Cox and Miller (1965) and Benjamin and Cornell (1970) can be mentioned. Markov chains, especially the Markov chains of the first-order and second-order, have been widely used in climatology for modeling the behaviour of wet and drought periods or transfer from a wet period to a dry one. Gabriel and Neumann (1962) were among the first people who used this model to study the features of daily precipitation occurrence. According to this model, they showed that daily precipitation in Tel Aviv has the features of two-state first-order Markov chain. In other words, there is a dependency between today's precipitation occurrence and yesterday's governing conditions. However, after the study of Gabriel and Neumann (1962), more complicated models were suggested to estimate the statistical distributions on meteorological data, but using this model has still a special place in climatology and meteorological studies. So that, Todorovic and Woolhieser (1975) used this model to study n-day precipitation and Hopkins and Robillard (1964) studied the daily precipitation of Prairie Province in Canada with this model. Victor and Sastry in

1979 studied the probability features of daily precipitation in Indian agricultural research institute by using this model for a 30-year time period from June to September. The results of this study show that daily precipitations of this station in the studied four months have the best fit at $\alpha = 0.01$ probability level with first-order Markov chain and at $\alpha = 0.05$ probability level with the second-order Markov chain for two months of June and September. Singh *et al.* (1984) also showed that the second-order Markov chain during the summer Monsoon of semi-dry regions in India has the best fit on daily precipitation data in this part of India. Thus, they used this model to study and analyze the probabilities of precipitation in this region. Subramanian and Sanjeeva (1989) are two other researchers who studied the occurrence of dry and wet monsoons of south Andhra Pradesh with the aim of achieving an appropriate crop calendar for dry farming by using the mentioned model. They used the weekly statistics of a 30-year period of precipitation from June to September. The intended threshold of these researchers to determine wet and dry weeks was 20 mm per week because they believed that this amount of precipitation can supply all the water requirements of the plants in then studied region during one week. First-order Markov chain model was the first model selected by these researchers to study the precipitation of this region. Other studies that used this model to identify the precipitation features of one region are the studies of Chin (1977), Jones and Thornton (1993), Steineman (2003), Moon *et al.* (1994), Martin-Vide and Gomez (1999), Lu & Berliner (1999) and Penal and Liano (2006).

Many studies have been carried out in Iran by using this model. They can be classified into two groups in terms of spatial scale: the first group includes the studies that examined the behaviour of wet and dry days in the daily scale (Saligheh *et al.*, 2012; Amini *et al.*, 2011; Asakereh, 2008; Asakereh and Mazini, 2010; Tavousi *et al.*, 2010; Banivahab, 2012; Jalali *et al.*, 2011) and the second group includes the studies that examined the behaviour of dry and wet behaviours in the monthly, seasonal and annual scales (Alizadeh and Tousi, 2008; Alizadeh *et al.*, 2008; Bashirzadeh and Araghi Nejad, 2010; Razei *et al.*, 2007; Alijani *et al.*, 2012). Mahmoudi *et al.* (2013) in a comprehensive study regionalized Iran according to the length of dry spells. They fitted Markov chains of different orders according to the length of dry spells with 0.1 mm threshold to show that at the studies with very long dry spells, the Markov chains with lower orders have the best fit, but at the stations that the length of dry spells is not long, the Markov chains with higher orders show the best fit.

In general, the review of available references on the use of Markov chain models in studying the spatial-




Fig. 1. The map of geographical location of the studied stations in Iran

temporal behaviour of dry and wet days showed that no comprehensive and complete study has been carried out in Iran to include the entire country as a whole unit. All of the previous studies that modeled the behaviour of wet and dry days or examined either one Meteorological station or one province. Thus, the purpose of this study is to model the behaviour of wet and dry days in Iran and extract some of its statistical features by using Markov chain models in order to develop the climatological knowledge of wet and dry days in Iran.

2. Data and methodology

To model the behaviour of wet and dry days in Iran and achieve a collection of appropriate patterns in this field, the data related to daily precipitation of 44 synoptic stations for a 25-year period (1991-2015) were extracted from Iran meteorological organization. These data were classified in 4 seasons of winter (December, January and February), spring (March, April and May), summer (June, July and August) and autumn (September, October and

November). Distribution and scattering of the studied stations in Iran were given in Fig. 1.

After extraction and formation of a database, the second step of the procedure was to determine a threshold to distinguish wet days from dry days. There is much disagreement among climatologists on determining this threshold, so that they suggested different criteria like 0.01, 0.1, 0.15, 0.2, 0.25 and 0.3 for this purpose (Domroes and Ranatung, 1993). Also, the World Meteorological Organization (WMO) defined a wet day with at least 1 mm of precipitation for 24 hours (Alijani *et al.*, 2014). In this study, according to environmental conditions of Iran and low number of wet days at some stations during the year, the value of 0.1 mm precipitation per day was determined as the threshold of a wet day. Now by determining the threshold and isolating days into two groups of dry days with precipitation threshold less than 0.1 mm ($X_t = 0$) and wet days with precipitation threshold more than 0.1 mm ($X_t = 1$), the data are ready for

fit on Markov chain. The most usual model of Markov chain is the first-order Markov chain that is defined as follows:

$$P_r \{X_{t+1} | X_t, X_{t-1}, \dots, X_1\} = P_r \{X_{t+1} | X_t\} \quad (1)$$

In this regard, P_r is conditional probability and indicates that the result of such process at $t+1$ depends only on conditions at time t . In other words, the prediction of tomorrow conditions is exclusively carried out by today's data and the yesterday's data do not provide any additional data. The first step to use the Markov chain model is to provide transition frequency matrix for calculating transition probability matrix. Transition frequency matrix for two-state first-order Markov chain is presented as follows:

$$N = \begin{matrix} & \begin{matrix} W & D \end{matrix} \\ \begin{matrix} W \\ D \end{matrix} & \begin{bmatrix} n_{00} & n_{01} \\ n_{10} & n_{11} \end{bmatrix} \end{matrix}$$

where, n_{ij} shows the frequency of transition states from state i to other possible states j (Moon *et al.*, 1994). In the next step, the transition probability matrix of stations must be calculated by using equation (2) (Alijani *et al.*, 2014).

$$P_{ij} = \frac{n_{ij}}{n_{i+}} = \frac{n_{ij}}{n_{i0} + n_{i1}} \quad (2)$$

This equation shows that if time series at time $t-1$ is at state i , the probability of the next value of time series as $X_{t+1}=j$ will be equal to P_{ij} . The matrix obtained from the results of this equation is called as transition probability matrix that is as follows:

$$P = \begin{matrix} & \begin{matrix} W & D \end{matrix} \\ \begin{matrix} W \\ D \end{matrix} & \begin{bmatrix} p_{00} & p_{01} \\ p_{10} & p_{11} \end{bmatrix} \end{matrix}$$

where, p_{ij} shows the transition probability from state i to other states j (Moon *et al.*, 1994). After providing transition frequency matrix and transition probability matrix, the independence test is used to study the dependent or independent wet and dry days. In the independence test, the null hypothesis is based on the idea that the series are independent of each other. That is, the data do not follow the first-order Markov this test is based on observed values of crosstab table (O) and expected values (E) under null hypothesis (Asakereh, 2008).

$$e_{ij} = \frac{n_i \times n_j}{\sum n} \quad (3)$$

where, n_i is the total rows of each frequency matrix and n_j is the total columns of each frequency matrix. Thus, finally the test statistic is obtained from the following equation:

$$x^2 = \sum_{i=1}^r \sum_{j=1}^c \frac{(O_{ij} - E_{ij})^2}{E_{ij}} \quad (4)$$

Also, the critical value is calculated based on $\alpha = 0.05$ and $df = (r - 1)(c - 1)$. Here, r and c are the rows and columns of matrix, respectively. If $x^2 > X_c$, the null hypothesis at the supposed significance level is rejected in which X_c is the corresponding critical value from chi square distribution with $(r - 1)(c - 1)$ degree of freedom.

Transition probability matrices make the calculation of different aspects of dry and wet days behaviour at each synoptic station possible in terms of different states and orders of Markov chain as follows (Lana and Burgueno, 1998):

The probabilities of detecting n consecutive dry days at a synoptic station according to first-order Markov chain include:

$$Q1(n) = (1 - P_{01})^{n-1} P_{01} \quad n \geq 1 \quad (5)$$

The probability of n -day state occurrence after a fixed starting state accuracy to the concept of two state first-order Markov chain matrix $P1_2$, can be calculated as follows:


$$S1_2(n) = V \cdot P1_2^n \quad (6)$$

That V will be the vector of components (0, 1), if the starting state is related to a wet day and it will be the vector of components (1, 0) if the starting state is a dry day.

A more interesting aspect is to determine starting values for the components of matrix $P1_2^n$ that will show the probabilities of stationary transition for different states of transition.

$$P1_2^n(0,0) = P1_2^n(1,0) = P_{10} / (P_{01} + P_{10}) \quad (7)$$

$$P1_2^n(0,1) = P1_2^n(1,1) = P_{01} / (P_{01} + P_{10}) \quad (8)$$



Figs. 2(a-d). Frequency percent of dry days for seasons in Iran. (a) Winter, (b) Spring, (c) Summer and (d) Autumn

These equations respectively show the columns of the first and second starting of matrix $P1_2^n$. The 2×2 starting matrix is then composed with 2 columns. The first column is the starting probability of a dry day at a synoptic station and second column is a wet day.

The components of transition matrix provide the context for the calculation of the expected length of dry or wet period in addition to the expected length of a dry-wet cycle. Thus, to calculate the length of wet-dry cycle, first the probabilities of detecting n consecutive wet or dry days must be calculated.

$$El(r) = 1 / P_{10} \quad (9)$$

$$El(d) = 1 / P_{01} \quad (10)$$

Then, the expected length of a dry-wet period can be calculated as follows:

$$El(c) = El(r) + El(d) = (P_{10} + P_{01}) / (P_{10}P_{01}) \quad (11)$$

Also, the probabilities to return to a dry day after n consecutive days, $ER(d)$, or one wet day after n consecutive days, $ER(r)$, can be stated as follows:

$$ER(r) = (P_{01} + P_{10}) / P_{01} \quad (12)$$

$$ER(d) = (P_{01} + P_{10}) / P_{10} \quad (13)$$

3. Results

The frequency percent of dry days was calculated for all 44 studied stations in 4 seasons and their results were given as 4 maps in Figs. 2(a-d). According to these maps, it is observed that the average percent of dry days on the south coast of the Caspian Sea is less than 70% in autumn and it reaches to more than 94% in Oman Sea coast in southeast Iran. In this season, Bandar Anzali with 52.9%

TABLE 1

The results of goodness of fit test to determine the conformity of two state first-order Markov chain with frequency of different states of dry and wet days. The homes with red color show the lack of conformity

Station	Winter	Spring	Summer	Autumn	Station	Winter	Spring	Summer	Autumn
Ahwaz	✓	✓	*	✓	Zahedan	✓	✓	*	✓
Abadan	✓	✓	✓	✓	Sanandaj	✓	✓	*	✓
Arak	✓	✓	✓	✓	Zanjan	✓	✓	✓	✓
Ardabil	✓	✓	✓	✓	Sabzevar	✓	✓	*	✓
Babolsar	✓	✓	✓	✓	Saghez	✓	✓	✓	✓
Bam	✓	✓	*	✓	Semnan	✓	✓	✓	✓
Bandar Abbas	✓	✓	*	✓	Shahrod	✓	✓	✓	✓
Bandar Anzali	✓	✓	✓	✓	Esfahan	✓	✓	✓	✓
Bandar Lengeh	✓	*	✓	✓	Shahre Kurd	✓	✓	*	✓
Boshehr	✓	✓	*	✓	Shiraz	✓	✓	*	✓
Birjand	✓	✓	*	✓	Tabas	✓	✓	*	✓
Tabriz	✓	✓	✓	✓	Hamedan	✓	✓	✓	✓
Torbat Hidaye	✓	✓	*	✓	Fasa	✓	✓	*	✓
Tehran	✓	✓	✓	✓	Ghazvin	✓	✓	✓	✓
Jask	✓	*	✓	*	Kashan	✓	✓	*	✓
Chabahar	✓	*	*	*	Kerman	✓	✓	*	✓
Khoram Abad	✓	✓	*	✓	Kermanshah	✓	✓	*	✓
Khoy	✓	✓	✓	✓	Gorgan	✓	✓	✓	✓
Dezful	✓	✓	*	✓	Mashhad	✓	✓	*	✓
Doshan Tapeh	✓	✓	✓	✓	Nozheh	✓	✓	✓	✓
Ramsar	✓	✓	✓	✓	Yazd	✓	✓	*	✓
Rasht	✓	✓	✓	✓	Oroomieh	✓	✓	*	✓
Zabol	✓	✓	*	✓	Iranshar	✓	✓	*	✓

and Jask with 96.9% had respectively the minimum and maximum frequency percent of dry days.

In winter that is considered as the wettest season in Iran, these values change slightly than autumn, so the south coast of the Caspian Sea in this season had less than 67% and south coasts of southeast Iran had an average more than 91%. The minimum and maximum frequency percent of dry days in this season belong to Rasht station with 55.5% and Jask station with 92.1% respectively.

In spring, the number of wet days in some parts of Iran significantly reduces and this reduction is clearly seen in the map related to spring in Figs. 2(a-d). In this season, Chabahar station with 99% in southeast Iran had the maximum frequency percent of dry days and Ardabil station in northwest Iran with 68.2% had the minimum frequency percent of dry days.

In summer, subtropical high pressure on the upper atmosphere of Iran, there is a very dry season in Iran, so that except the northern coast of Iran, the whole country has more than 90% dry days and at Zabol station, 100% of the studied days were dry.

To model the behaviour of dry and wet days in Iran, the first-order Markov chain was used. Thus, for all 44 studied stations, first transition frequency matrices and then transition probability matrices were adjusted. The number of matrices prepared at this step included 352 matrices; 176 matrices were related to transition frequency matrix and 176 matrices were related to transition probability matrix. After the adjustment of matrices, the chi-square test χ^2 was used for goodness of fit of first-order Markov chain at the frequency of different states of wet and dry days at stations. The obtained results were given in Table 1. In this table, the seasons that the

TABLE 2

The probabilities of occurring 5, 10, 15, 20, 25 and 30 dry periods in autumn at synoptic stations of Iran

Station	5 days	10 days	15 days	20 days	25 days	30 days	Station	5 days	10 days	15 days	20 days	25 days	30 days
Ahwaz	0.07	0.04	0.02	0.01	0.01	0	Zahedan	0.02	0.02	0.02	0.02	0.01	0.01
Abadan	0.06	0.04	0.02	0.01	0.01	0.1	Sanandaj	0.08	0.03	0.02	0.01	0	0
Arak	0.08	0.04	0.02	0.1	0	0	Zanjan	0.08	0.03	0.01	0.01	0	0
Ardabil	0.08	0.03	0.01	0.01	0	0	Sabzevar	0.07	0.04	0.02	0.01	0.01	0
Babolsar	0.08	0.02	0.01	0	0	0	Saghez	0.08	0.03	0.01	0	0	0
Bam	0.03	0.02	0.02	0.02	0.01	0.01	Semnan	0.06	0.04	0.02	0.02	0.01	0.01
Bandar Abbas	0.03	0.02	0.02	0.02	0.01	0.01	Shahrod	0.07	0.04	0.02	0.01	0.01	0
Bandar Anzali	0.08	0.02	0	0	0	0	Esfahan	0.06	0.04	0.02	0.02	0.01	0.01
Bandar Lengeh	0.03	0.03	0.02	0.02	0.02	0.01	Shahre Kurd	0.07	0.04	0.02	0.01	0.01	0
Boshehr	0.06	0.04	0.02	0.02	0.01	0.01	Shiraz	0.06	0.04	0.02	0.01	0.01	0.01
Birjand	0.05	0.04	0.03	0.02	0.01	0.01	Tabas	0.05	0.03	0.03	0.02	0.01	0.01
Tabriz	0.08	0.03	0.01	0.01	0	0	Hamedan	0.08	0.04	0.02	0.01	0	0
Torbat Hay	0.06	0.04	0.02	0.01	0.01	0.01	Fasa	0.05	0.04	0.03	0.02	0.01	0.01
Tehran	0.07	0.04	0.02	0.01	0	0	Ghazvin	0.08	0.03	0.02	0.01	0	0
Jask	0.02	0.02	0.02	0.01	0.01	0.01	Kashan	0.06	0.04	0.02	0.02	0.01	0.01
Chabahar	0.02	0.02	0.02	0.01	0.01	0.01	Kerman	0.05	0.04	0.03	0.02	0.01	0.01
Khoram Abad	0.08	0.04	0.02	0.01	0	0	Kermanshah	0.08	0.04	0.02	0.01	0	0
Khoy	0.08	0.03	0.02	0.01	0	0	Gorgan	0.08	0.03	0.01	0	0	0
Dezful	0.07	0.04	0.02	0.01	0.01	0	Mashhad	0.07	0.04	0.02	0.01	0.01	0
Doshan Tapeh	0.07	0.04	0.02	0.01	0	0	Nozheh	0.08	0.03	0.02	0.01	0	0
Ramsar	0.08	0.02	0	0	0	0	Yazd	0.02	0.03	0.02	0.02	0.02	0.01
Rasht	0.08	0.02	0	0	0	0	Oroomich	0.08	0.04	0.02	0.01	0	0
Zabol	0.03	0.02	0.02	0.02	0.01	0.01	Iranshhar	0.02	0.02	0.02	0.02	0.01	0.01

frequency of their wet and dry days was not consistent with first-order Markov chain, *i.e.*, their dry and wet days were independent from each other, were shown with red color. In this table, it is observed that in summer, the first-order Markov chain was not consistent with most stations because there were a few wet days but in winter, due to the high number of wet days that first-order Markov chain is consistent with the frequency of different states of dry and wet days at stations.

The other reason that only first-order Markov chain was used in this study and Markov chains with higher orders were not used was due to two reasons. Firstly, the first-order Markov chain except summer showed the best conformity with frequency of different states of dry and wet days at most stations in three other seasons. The second reason was related to limited number of databases

at stations because the small size of databases on the one hand and high number of states related to wet and dry days at higher orders of Markov chain makes some mistakes in calculation of probabilities. Thus, here the Markov features of different states of wet and dry days in Iran are explained according to two- state first-order Markov chain.

3.1. The probabilities of detecting n consecutive dry days

The possibility of occurrence of dry day's continuity of 5, 10, 15, 20, 25 and 30-days was the first Markov feature that was calculated using Equation 5 for all stations studied in different seasons of the year. The results of the calculation of the Markov feature have been presented in Tables (2-4) for autumn, winter and spring,

TABLE 3


The probabilities of occurring 5, 10, 15, 20, 25 and 30 dry periods in winter at synoptic stations of Iran

Station	5 days	10 days	15 days	20 days	25 days	30 days	Station	5 days	10 days	15 days	20 days	25 days	30 days
Ahwaz	0.08	0.03	0.01	0	0	0	Zahedan	0.08	0.04	0.02	0.01	0.01	0
Abadan	0.08	0.04	0.02	0.01	0	0	Sanandaj	0.08	0.02	0	0	0	0
Arak	0.08	0.02	0.01	0	0	0	Zanjan	0.08	0.02	0	0	0	0
Ardabil	0.06	0.04	0.02	0.02	0.01	0.01	Sabzevar	0.08	0.03	0.01	0	0	0
Babolsar	0.08	0.02	0.01	0	0	0	Saghez	0.08	0.02	0	0	0	0
Bam	0.06	0.04	0.02	0.02	0.01	0.01	Semnan	0.08	0.04	0.02	0.01	0	0
Bandar Abbas	0.07	0.04	0.02	0.01	0.01	0	Shahrod	0.08	0.03	0.02	0.01	0	0
Bandar Anzali	0.08	0.02	0	0	0	0	Esfahan	0.08	0.04	0.02	0.01	0	0
Bandar Lengeh	0.07	0.04	0.02	0.01	0.01	0	Shahre Kurd	0.08	0.02	0.01	0	0	0
Boshehr	0.08	0.03	0.01	0	0	0	Shiraz	0.08	0.03	0.01	0	0	0
Birjand	0.08	0.03	0.01	0	0	0	Tabas	0.08	0.04	0.02	0.01	0	0
Tabriz	0.08	0.03	0.01	0	0	0	Hamedan	0.08	0.02	0	0	0	0
Torbat Hay	0.08	0.03	0.01	0	0	0	Fasa	0.08	0.03	0.02	0.01	0	0
Tehran	0.08	0.03	0.01	0	0	0	Ghazvin	0.08	0.02	0.01	0	0	0
Jask	0.06	0.04	0.02	0.02	0.01	0.01	Kashan	0.08	0.03	0.02	0.01	0	0
Chabahar	0.05	0.03	0.03	0.02	0.01	0.01	Kerman	0.08	0.03	0.01	0.01	0	0
Khoram Abad	0.08	0.02	0	0	0	0	Kermanshah	0.08	0.02	0	0	0	0
Khoy	0.08	0.03	0.01	0.01	0	0	Gorgan	0.08	0.02	0	0	0	0
Dezful	0.08	0.03	0.01	0	0	0	Mashhad	0.08	0.02	0.01	0	0	0
Doshan Tapeh	0.08	0.03	0.01	0	0	0	Nozheh	0.08	0.02	0	0	0	0
Ramsar	0.08	0.02	0.01	0	0	0	Yazd	0.08	0.03	0.02	0.01	0	0
Rasht	0.08	0.03	0.01	0	0	0	Oroomieh	0.08	0.03	0.01	0	0	0
Zabol	0.02	0.02	0.02	0.01	0.01	0.01	Iranshhar	0.02	0.02	0.02	0.01	0.01	0.01

TABLE 4

The probabilities of occurring 5, 10, 15, 20, 25 and 30 dry periods in spring at synoptic stations of Iran

Station	5 days	10 days	15 days	20 days	25 days	30 days	Station	5 days	10 days	15 days	20 days	25 days	30 days
Ahwaz	0.03	0.02	0.02	0.02	0.02	0.01	Zahedan	0.03	0.03	0.02	0.02	0.02	0.01
Abadan	0.03	0.03	0.02	0.02	0.02	0.01	Sanandaj	0.07	0.04	0.02	0.01	0	0
Arak	0.07	0.04	0.02	0.01	0.01	0	Zanjan	0.08	0.03	0.01	0	0	0
Ardabil	0.08	0.02	0.01	0	0	0	Sabzevar	0.07	0.04	0.02	0.01	0.01	0
Babolsar	0.08	0.04	0.02	0.01	0	0	Saghez	0.08	0.03	0.02	0.01	0	0
Bam	0.03	0.03	0.02	0.02	0.02	0.01	Semnan	0.06	0.04	0.02	0.01	0.01	0.01
Bandar Abbas	0.02	0.02	0.02	0.01	0.01	0.01	Shahrod	0.08	0.04	0.02	0.01	0	0
Bandar Anzali	0.08	0.02	0.01	0	0	0	Esfahan	0.05	0.03	0.03	0.02	0.01	0.01
Bandar Lengeh	0.02	0.02	0.02	0.01	0.01	0.01	Shahre Kurd	0.06	0.04	0.02	0.01	0.01	0.1
Boshehr	0.03	0.02	0.02	0.02	0.01	0.01	Shiraz	0.05	0.03	0.03	0.02	0.01	0.1
Birjand	0.05	0.03	0.03	0.02	0.01	0.01	Tabas	0.03	0.03	0.02	0.02	0.02	0.1
Tabriz	0.08	0.03	0.01	0	0	0	Hamedan	0.08	0.04	0.02	0.01	0	0
Torbat Hay	0.07	0.04	0.02	0.01	0.01	0	Fasa	0.03	0.02	0.02	0.02	0.01	0.01
Tehran	0.07	0.04	0.02	0.01	0	0	Ghazvin	0.08	0.03	0.01	0	0	0
Jask	0.02	0.02	0.02	0.01	0.01	0.01	Kashan	0.05	0.04	0.03	0.02	0.01	0.01
Chabahar	0.01	0.01	0.01	0.01	0.01	0.01	Kerman	0.04	0.03	0.02	0.02	0.01	0.01
Khoram Abad	0.07	0.04	0.02	0.01	0.01	0	Kermanshah	0.07	0.04	0.02	0.01	0	0
Khoy	0.08	0.02	0.01	0	0	0	Gorgan	0.08	0.03	0.01	0	0	0
Dezful	0.05	0.03	0.03	0.02	0.01	0.01	Mashhad	0.08	0.03	0.02	0.01	0	0
Doshan Tapeh	0.07	0.04	0.02	0.01	0.01	0	Nozheh	0.08	0.03	0.02	0.01	0	0
Ramsar	0.08	0.02	0.01	0	0	0	Yazd	0.08	0.03	0.02	0.01	0	0
Rasht	0.08	0.03	0.01	0	0	0	Oroomieh	0.08	0.03	0.01	0	0	0
Zabol	0.02	0.02	0.02	0.01	0.01	0.01	Iranshhar	0.02	0.02	0.02	0.01	0.01	0.01



Figs. 3(a-c). Expected dry spells length for different seasons of Iran (in terms of day). (a) Autumn, (b) Winter and (c) Spring

respectively. According to the Tables (2-4), the possibility of occurrence of each of the mentioned continuities (5, 10, 15, 20, 25 and 30 days) was calculated for each station and season. For instance, the possibility of occurrence of a 5-days dry period in autumn is 0.07 in Ahwaz station and/or the possibility of occurrence of a 30-days period in Zahedan station is 0.01 (Table 2).

According to the results of this Markov feature, it is observed that there is the probability of occurring 5 and 10 dry consecutives in 3 seasons of autumn, winter and spring for all studied stations with different probabilities [Tables (2-4)]. But for consecutives higher than 15 days, it is observed that at some stations and seasons, their probability of occurring becomes zero. For example, 3 stations of Bandar Anzali, Ramsar and Rasht do not have dry periods more than 15 days due to their precipitation regime in autumn (Table 2).


In winter, with the arrival of wet waves to Iran and since the precipitation regime of most stations in Iran are winter, the probability of occurring long term consecutive dry days is very low, so that a large number of stations like Bandar Anzali, Ramsar, Rasht and Gorgan at southern coasts of Caspian sea and Khorram Abad stations, Sanandaj, Saghez, Zanjan, Hamedan and Kermanshah in western Iran do not have the probability of occurring more than 15 consecutive days [Figs. 3(a-c)].

In spring, all stations experience 5, 10 and 25 day consecutives with different probabilities from 0.01 to 0.08. In addition, the number of stations that have 30-day consecutives has increased as can be seen in Table 4.

In contrast, the results for some stations such as Bandar Lengeh, Jask and Chabahar in autumn and spring are very much different from other stations. Considering Tables (2&4), it can be seen that the possibility of

TABLE 5
Stationary transition probability to a dry day with wet day at synoptic stations of Iran

Station	Stationary transition probability to a dry day or a wet day					
	Autumn		Winter		Spring	
	Transition probability to a dry day	Transition probability to a wet day	Transition probability to a dry day	Transition probability to a wet day	Transition probability to a dry day	Transition probability to a wet day
Ahwaz	0.86	0.14	0.77	0.23	0.94	0.06
Abadan	0.87	0.13	0.92	0.18	0.95	0.05
Arak	0.79	0.21	0.66	0.34	0.82	0.18
Ardabil	0.77	0.23	0.87	0.13	0.68	0.32
Babolsar	0.67	0.33	0.66	0.34	0.81	0.19
Bam	0.96	0.04	0.89	0.11	0.95	0.05
Bandar Abbas	0.96	0.04	0.85	0.15	0.98	0.02
Bandar Anzali	0.53	0.47	0.56	0.44	0.71	0.29
Bandar Lengeh	0.94	0.06	0.86	0.14	0.97	0.03
Boshehr	0.87	0.13	0.76	0.24	0.96	0.04
Birjand	0.90	0.10	0.75	0.25	0.91	0.09
Tabriz	0.77	0.23	0.71	0.29	0.71	0.29
Torbat Hay	0.86	0.14	0.69	0.31	0.85	0.15
Tehran	0.79	0.21	0.73	0.27	0.83	0.17
Jask	0.96	0.04	0.89	0.11	0.97	0.03
Chabahar	0.97	0.03	0.92	0.08	0.99	0.01
Khoram Abad	0.77	0.23	0.64	0.36	0.83	0.17
Khoy	0.79	0.21	0.78	0.22	0.68	0.32
Dezful	0.83	0.17	0.72	0.28	0.92	0.08
Doshan	0.80	0.20	0.72	0.28	0.83	0.17
Ramsar	0.64	0.36	0.63	0.37	0.68	0.32
Rasht	0.58	0.42	0.55	0.45	0.69	0.31
Zabol	0.96	0.04	0.84	0.16	0.97	0.03
Zahedan	0.96	0.04	0.87	0.13	0.95	0.05
Sanandaj	0.76	0.24	0.65	0.35	0.81	0.19
Zanjan	0.75	0.25	0.66	0.34	0.72	0.28
Sabzevar	0.85	0.15	0.73	0.27	0.85	0.15
Saghez	0.73	0.27	0.63	0.37	0.77	0.23
Semnan	0.89	0.11	0.83	0.17	0.88	0.12
Shahrod	0.87	0.13	0.81	0.19	0.82	0.18
Esfahan	0.89	0.11	0.83	0.17	0.91	0.09
Shahre Kurd	0.81	0.19	0.70	0.30	0.87	0.13
Shiraz	0.86	0.14	0.71	0.29	0.91	0.09
Tabas	0.92	0.08	0.82	0.18	0.94	0.06
Hamedan	0.78	0.22	0.66	0.34	0.79	0.21
Fasa	0.88	0.12	0.78	0.22	0.95	0.05
Ghazvin	0.77	0.23	0.69	0.31	0.77	0.23
Kashan	0.89	0.11	0.81	0.19	0.90	0.10
Kerman	0.90	0.10	0.77	0.23	0.92	0.08
Kermanshah	0.75	0.25	0.63	0.37	0.80	0.20
Gorgan	0.73	0.27	0.64	0.36	0.74	0.26
Mashhad	0.84	0.16	0.66	0.34	0.78	0.22
Nozheh	0.76	0.24	0.66	0.34	0.77	0.23
Yazd	0.94	0.06	0.86	0.14	0.83	0.17
Oroomieh	0.78	0.22	0.73	0.27	0.75	0.25
Iranshahr	0.96	0.04	0.88	0.12	0.98	0.02



Figs. 4(a-c). Expected wet spells length for different seasons in Iran (in terms of day). (a) Autumn, (b) Winter and (c) Spring


occurrence of dry day's continuity of 5-30 days is almost the same for this station. The reasons for this similarity can be the dry climate of these stations, their winter precipitation regime and lack of Markov chain fit on the dry and wet days of these stations (Table 1). The same conditions can be seen in other stations such as Bam, Bushehr, Bandar Abbas, Bandar Lengeh, Zabol and Iranshahr except that the result of Chi-square test confirmed the Markov chain fit on their dry and wet days. (Table 1).

3.2. Stationary transition probabilities of dry and wet days occurrence

By multiplying transition probability matrix, a situation is occurred in which all rows of transition probability matrix become equal to each other, so that if the process of multiplication continues after this, no

change will be observed in transition probability matrix arrays. Such a matrix is called stationary transition probability matrix. This matrix is calculated for all studied stations by using equations 7 and 8 and the results were observed in Table 5 for 3 seasons of autumn, winter and spring.

As this table shows, the long term transition probability of the studied situations from one state to another one. For example, stationary transition probability matrix of autumn in Ahwaz station shows that transition probability to one dry day is equal to 0.86 and to a wet day is equal to 0.14. The lowest transition probability to a dry day in Iran in autumn belongs to Bandar Anzali station, in winter belongs to Rasht station and in spring belongs to Ardabil, Khoy and Ramsar stations. Certainly, the maximum transition probability to a dry day in all 3 seasons belongs to Chabahar station in southeast



Figs. 5(a-c). Expected wet-dry cycle length for different seasons in Iran (in terms of day). (a) Autumn, (b) Winter and (c) Spring

Iran that is respectively equal to 0.97, 0.92 and 0.99 for autumn, winter and spring.


3.3. Dry-wet spells cycle

Another important feature that can be extracted from transition probability matrix of stations for understanding the wet and dry days' behaviour in Iran is the expected dry and wet spells length. Expected dry and wet spells length is in fact the long term average (mathematical expectation) of wet and dry spells lengths that was calculated by equations 9 and 10 for all studied stations. Their results were given in the form of contour maps [Figs. (3&4)]

As Figs. 3(a-c) shows, the expected dry and wet spells length shows different behaviours in different seasons of Iran. so that, the difference between minimum and maximum expected dry spells length between the

northern part and southern part of Iran in autumn is relatively high (Bandar Anzali 3.7 days and Chabahar 50 days). However, this difference reaches to its minimum amount in winter so that the difference between the northern and southern parts of Iran reaches to 12 days in this season (Rasht 3.6 days and Chabahar 16.7 days). In spring, since it is a transition spring, the difference between expected dry days length in the country increases again [Figs. 3(a-c)].

Expected wet spells length has a more equal behaviour in comparison to expected dry spells length. In other words, there is not much difference between the expected wet spells in the northern and southern parts of Iran. In autumn, the difference between expected wet spells length between the two stations has the maximum and minimum values if almost 1.6 days (Rasht = 2.9 days and Iranshahr = 1.3 day). In winter, this value reaches to 1.4 days (Rasht = 2.9 days and Semnan = 1.5 day) and in



Figs. 6(a-c). Return periods for dry days for different seasons in Iran (in terms of day). (a) Autumn, (b) Winter and (c) Spring

spring to 1 day (Rasht = 1.2 day and Iranshahr = 1.1 day) [Figs. 4(a-c)]

From the total values of expected dry-wet spells length, the expected dry-wet cycle for each station is obtained. This feature of Markov chain was calculated by equation 11 for all studied stations and its results were given in the form of 3 maps in Figs. 5(a-c).


According to Figs. 5(a-c), it is observed that wet-dry spells length in winter is not very long for the entire Iran and this is due to the climatic status of this country in this season. The shortest wet-dry cycle length in this season belongs to Gorgan station with 5.7 days and its longest length belongs to Chabahar station with 8.2 days. In autumn, the difference between maximum and minimum dry-wet cycle length between the southern and northern parts of Iran increases (Ramsar = 6.3 days and

Jask = 51.9 days) and in spring, this difference reaches to its maximum level (Ardabil = 8.1 days and Chabahar = 101.3 days).

3.4. Return periods for dry or wet episode

Figs. (6&7) show the return periods obtained for a new dry or wet day. In Figs. 6(a-c), as can be seen, the expected return periods for dry days is almost fixed for the entire Iran, so that in autumn, it is between 1.03 to 1.87 days, in winter is between 1.09 to 1.82 days and in spring is between 1.01 to 1.46. Thus, it is concluded that there cannot be a significant difference between different parts of Iran.

But the number of return days to a new precipitation period for autumn is between 2.15 to 33.5 days, for winter is between 2.21 to 12 and in spring is



Figs. 7(a-c). Return periods for wet days for different seasons in Iran (in terms of day). (a) Autumn, (b) Winter and (c) Spring

between 3.17 to 79 days that emphasizes the climatic diversity of Iran [Figs. 7(a-c)].

4. Conclusions

The behaviour of dry and wet days in Iran was modeled according to Markov chains. For this modeling, the transition frequency matrix of first-order Markov chain was adjusted for all studied stations, for goodness of fit test to determine the appropriateness of first-order Markov chain, the chi square test was used. The results of this test showed that different states of wet and dry consecutives in Iran in summer had no Markov behaviour. Thus, summer was excluded from calculations. Then, according to the conformity of two - state first-order Markov chain at different states of wet and dry days, it was made a decision to select first-order Markov chain for extraction of behavioural features of dry and wet days in Iran.

The average frequency percent of dry days in Iran shows that in winter that is considered as the wet season in Iran, changes about 55.5 % in northern Iran to 91% in southeast Iran. This value varies from 52.9 to 96.9% in autumn, from 68.2 to 99.1 % in spring and from 90 to 100% in summer.

There is the occurrence of 5-10 dry days' consecutives in 3 seasons of autumn, winter and spring with different probabilities for the whole country. But the probability of occurring dry consecutives more than 20 days is very variable according to season and place. So that, in winter with the arrival of western waves to Iran, we see the lack of occurrence of these consecutives more than 20 days in the half north of Iran. In two seasons of spring and autumn that are the end and start of long term sustainable conditions in the upper atmosphere of Iran, the probability of occurrence of 30-day dry spells increases very high especially in the southern part of Iran.

The lowest transition probability to one dry day in autumn belongs to Bandar Anzali station, in winter belongs to Rasht station and in spring belongs to Ardabil, Khoy and Ramsar stations that show the wet state of these stations in Iran. But the important point is that this Markov chain feature is related to two stations of northwest Iran, *i.e.*, Ardabil and Khoy stations in spring. The reason that these two stations in this season have the lowest transition probability to a dry day is hidden in the precipitation type of this season in this region because apparently the most precipitations of this region in this season are of convective type especially in comparison to southern coast of Caspian Sea.

Expected return periods for dry days is almost constant throughout Iran and is in a range from 1 to 2 days but the number of return days to a precipitation period does not follow this rule and is very variable for the whole country, so that it varies from 2.15 days in autumn to 79 days in spring that represents the climatic diversity of Iran.

Finally, it can be concluded that the two-state first-order Markov chain model except summer can model the behaviour of dry and wet days in Iran and reveal most of its hidden features.

Acknowledgments

The authors would like to thank Iran Meteorological Organization for its providing of daily precipitation data of 44 synoptic stations for a 25-year time span. Also it should be noted that the contents and views expressed in this research paper are the views of the authors and do not necessarily reflect the views of the organizations they belong to.

References

- Alijani, B., Mahmoudi, P., Chougan, A. and Bishe Niyasar, M., 2012, "Review the Structure for Continuity of the Two States Annual Precipitation in South Part of Iran, by Using Latent State of Markov Chain", *Geography and Development Iranian Journal*, **9**, 25, 1-16 (In Persian).
- Alijani, B., Mahmoudi, P., Shahoozchi, A. and Mohammadi, A., 2014, "Study of the persistence of precipitation days in Iran", *Geography and Environmental Planning*, **25**, 4, 1-16. (In Persian)
- Alizadeh, A. and Tousi, S. A., 2008, "Development of a Model for Monitoring and forecasting drought (case study: Khorasan Razavi province)", *Journal of Water and Soil (Agricultural Sciences and Technology)*, **22**, 1, 223-234 (In Persian).
- Alizadeh, A., Ansari, H., Ershadi, S. and Tousi, S. A., 2008, "Drought predictability in the province of Sistan and Baluchestan", *Journal of Geography and Regional Development*, **11**, 1-19 (In Persian).
- Amini, E., Ghahraman, B., Davary, K. and Mousavi Baygi, M., 2011, "Estimation of the Daily Rainfall Amount in Province of Khorasan", *Journal of Water and Soil, (Agricultural Sciences and Technology)*, **25**, 5, 1147-1157, doi: 10.22067/jsw.v0i--1.1228 (In Persian).
- Asakereh, H. and Mazini, F., 2010, "Investigation of dry days occurrence probability in Golestan province using Markov chain model", *Geography and Development Iranian Journal*, **8**, 17, 29-44.
- Asakereh, H., 2008, "Analysis of the frequency and the spell of rainy days using Markov chain model for city of Tabriz, Iran", *Iran-Water Resources Research*, **4**, 2, 46-56 (In Persian).
- Babaie, O. and Farajzadeh, M., 2002, "Patterns of spatial and temporal variations of rainfall in Iran", *Quarterly MODARES Human Sciences*, **6**, 4, 51-76 (In Persian).
- Banivahab, A., 2012, "The analysis and forecasting of climatic fluctuation of khorasan", *Territory*, **8**, 1, 93-105 (In Persian).
- Bashirzadeh, M. and Araghi Nejad, S., 2010, "Forecasting severity, duration and frequency of droughts using Markov chain and run theories (Case study: Lorestan Province)", *Iranian Water Research Journal*, **4**, 6, 91-95 (In Persian).
- Benjamin, J. R. and Cornell, C. A., 1970, "Probability, statistics and decision for civil engineers", McGraw-Hill, Inc., New York.
- Burgueno, A., Codina, B., Redano, A. and Lorente, J., 1994, "Basic statistical characteristics of hourly rainfall amounts in Barcelona (Spain)", *Theoretical and Applied Climatology*, **9**, 175-181, doi: 10.1007/BF00865532.
- Chin, E., 1977, "Modeling daily precipitation occurrence process with Markov chain", *Water Resources Research*, **13**, 6, 949-956, doi: 10.1029/WR013i006p00949.
- Cox, D. R. and Miller, H. D., 1965, "The theory of Stochastic process", Wiley, New York.
- De Arruda, H. V. and Pinto, H. S., 1980, "An alternative model for dry-spell probability analysis", *Monthly Weather Review*, **108**, 6, 823-825, doi:http://dx.doi.org/10.1175/1520-0493(1980)108<0823:AAMFDS>2.0.CO;2.
- Domroes, M. and Ranatung, E., 1993, "A Statistical approach toward a regionalization of daily rainfall in Sri Lanka", *International Journal of Climatology*, **13**, 7, 741-754, doi: 10.1002/joc.337.0130704.
- Gabriel, K. R. and Neumann, J., 1962, "A Markov chain model for daily rainfall occurrence at Tel Aviv", *Quarterly Journal of the Royal Meteorological Society*, **88**, 375, 90-95, doi: 10.1002/qj.49708837511.
- Hopkins, J. W. and Robillard, P., 1964, "Some statistics of daily rainfall occurrence for the Canadian Prairie Province", *Journal of Applied Meteorology and Climatology*, **3**, 600-602, doi: http://dx.doi.org/10.1175/1520-0450(1964)003<0600:SSO DRO>2.0.CO;2.
- Jalali, M., Karegar, H. and Soltani, S., 2011, "Analysis of rainy days occurrence probability using Markov chain model for city of Urmia Iran", *Geographic Space*, **11**, 35, 235-257 (In Persian).
- Jones, P. G. and Thornton, P. K., 1993, "A rainfall generator for agriculture applications in the tropics", *Agricultural and Forest Meteorology*, **63**, 1-2, 1-19, doi: http://dx.doi.org/10.1016/0168-1923(93)90019-E.
- Kemeny, J. G. and Snell, J. L., 1960, "Finite Markov chains", Van Nostrand Co. Inc., London.
- Lana, X. and Burgueno, A., 1998, "Daily dry-wet behavior in Catalonia (NE Spain) from the viewpoint of Markov Chains", *International Journal of Climatology*, **18**, 7, 793-815, doi: 10.1002/(SICI)1097-0088(19980615)18:7<793::AID-JOC262>3.0.CO;2-M.

- Lu, Z. Q. and Berliner, L. M., 1999, "Markov switching time series models with application to a daily runoff series", *Water Resources Research*, **35**, 2, 523-534, doi: 10.1029/98WR02686.
- Mahmoudi, P., Parvin, N. and Rezaei, J., 2013, "Regionalization of Iran based on the length of dry spells", *Arid Regions Geographic Studies*, **4**, 13, 85-106 (In Persian).
- Martin-Vide, J. and Gomez, L., 1999, "Regionalization of Peninsular Spain based on the length of dry spells", *International Journal of Climatology*, **19**, 5, 537-555, doi: 10.1002/(SICI)1097-0088(199904)19:5<537::AID-JOC371>3.0.CO;2-X.
- Moon, S. E., Ryoo, S. B. and Kwon, J. G., 1994, "A Markov chain model for daily precipitation occurrence in South Korea", *International Journal of Climatology*, **14**, 9, 1009-1016, doi: 10.1002/joc.3370140906.
- Movahedi, S., Asakereh, H., Sabziparvar, A. A., Masoodian, A. and Maryanaji, Z., 2012, "Assessment of Variability of Precipitation Regime in Iran", *Journal of Water and Soil*, **25**, 6, 1434-1447, doi: 10.22067/jsw.v0i0.12487 (In Persian).
- Nazaripour, H., 2011, "Synoptic Analysis of Iranian Rainfall Persistency", Ph. D. Thesis, University of Sistan and Baluchestan, Iran (In Persian).
- Penal, O. C. and Llano, M. P., 2006, "Temporal variability in the length of no-rain spells in Argentina", Proceedings of 8 ICSHMO, Foz do Iguaco, Brazil, April 24-28.
- Raziei, T., Daneshkar Arasteh, P., Akhtari, R. and Saghafian, B., 2007, "Investigation of meteorological droughts in the Sistan and Balouchestan province, using the standardized precipitation index and Markov chain model", *Iran-Water Resources Research*, **3**, 1, 25-35 (In Persian).
- Saligheh, M., Alijani, B. and Delara, G., 2012, "Spatial analysis of rainfall in the wet season, using Markov chain model case study: Ardabil province", *Journal of Geographical Sciences*, **20**, 23, 25-44 (In Persian).
- Singh, S. V., Kripalani, R. H., Saha, P. and Prasad, K. D., 1984, "Analysis and prediction of short period droughts during summer monsoon over India", *Mausam*, **35**, 361-366.
- Steinemann, A., 2003, "Drought indicators and triggers: A Stochastic approach to evaluation", *Journal of the American Water Resources Association*, **39**, 5, 1217-1233, doi: 10.1111/j.1752-1688.2003.tb03704.x.
- Subramaniam, A. R. and Sanjeeva, R. P., 1989, "Dry spell sequences in south coastal Andhra", *Mausam*, **40**, 57-60.
- Tavousi, T., Khosravi, M. and Ghaderi Zeh, K., 2010, "Assessment of drought and trends analysis of short-duration dry periods in Iranshahr region, using a Markov chain (1980-2006)", *Environmental Sciences*, **7**, 4, 31-44.
- Todorovic, P. and Woolhieser, D. A., 1975, "A Stochastic model of n-day precipitation", *Journal of Applied Meteorology*, **14**, 1, 17-24, doi: [http://dx.doi.org/10.1175/1520-0450\(1975\)014<0017:ASMODP>2.0.CO;2](http://dx.doi.org/10.1175/1520-0450(1975)014<0017:ASMODP>2.0.CO;2).
- Victor, U. S. and Sastry, P. S., 1979, "Dry spell probability by Markov chain model and its application to crop development stages", *Mausam*, **30**, 479-484.
- Zolfaghari, H., Hashemi, R. and Fashi, M., 2009, "Investigation of proportion of Daily Maximum precipitation to Annual precipitation in Iran", *Geographical Research*, **24**, 1, 167-188 (In Persian).