

A simple local variational iteration method for solving nonlinear Lane-Emden problems

Asghar Ghorbani^{a,*}, Mojtaba Bakherad^b

^a*Department of Applied Mathematics, Faculty of Mathematical Sciences, Ferdowsi University of Mashhad, Mashhad, Iran*

^b*Department of Mathematics, Faculty of Mathematics, University of Sistan and Baluchestan, Zahedan, Iran*

Abstract

In this paper, an explicit analytical method called the variational iteration method (VIM) is presented for solving the second-order singular initial value problems of Lane-Emden type, and its local convergence is discussed. Since it is often useful to have an approximate analytical solution to describe the Lane-Emden type equations, especially for ones where the closed-form solutions do not exist at all, therefore an effective improvement of the VIM is further proposed that is capable of obtaining an approximate analytical solution. The improved VIM is then treated as a local algorithm in a sequence of intervals as well as an adaptive one for finding accurate approximate solutions of the nonlinear Lane-Emden type equations. Some examples are given to demonstrate the efficiency and accuracy of the proposed method.

Keywords: Local variational iteration method, Truncated method, Adaptive strategy, Lane-Emden equations

1. Introduction

Recently, a lot of attention has been focused on the study of singular initial value problems (IVPs) in the second-order ordinary differential equations

*Corresponding author

Email address: aghorbani@um.ac.ir (Asghar Ghorbani)

(ODEs). Many problems in mathematical physics and astrophysics can be modeled by the so-called IVPs of the Lane-Emden type equation [2, 4, 15]:

$$\begin{cases} y'' + \frac{2}{x}y' + f(x, y) = g(x), \\ y(0) = a, \quad y'(0) = b, \end{cases} \quad (1)$$

where a and b are constants, $f(x, y)$ is a continuous real valued function, and $g(x) \in C[0, \infty]$. When $f(x, y) = K(y)$, $g(x) = 0$, Eq. (1) reduces to the classical Lane-Emden equation which, with specified initial conditions, was used to model several phenomena in mathematical physics and astrophysics such as the theory of stellar structure, the thermal behavior of a spherical cloud of gas, isothermal gas sphere, and the theory of thermionic currents [2, 4, 15].

Since the Lane-Emden type equations have significant applications in many fields of scientific and technical world, a variety of forms of $f(x, y)$ and $g(x)$ have been investigated by many researchers (e.g., [3, 16, 17]). A discussion of the formulation of these models and the physical structure of the solutions can be found in the literature. Though the numerical solution of the Lane-Emden equation (1), as well as other various linear and nonlinear singular IVPs in quantum mechanics and astrophysics [9], is numerically challenging because of the singularity behavior at the origin $x = 0$, but analytical solutions are much needed for physical understanding. Recently, many analytical methods were used to solve Lane-Emden equation [8, 10, 18]. Those methods are based on either series solutions or perturbation techniques [1, 11, 13, 14]. However, the convergence region of the corresponding results is very small.

The strategy that will be pursued in this work rests mainly on establishing a useful algorithm based on the variational iteration method (VIM) [7, 5] for finding highly accurate solution of the Lane-Emden type equations that it

- Overcomes main difficulty arising in the singularity of the equation at $x = 0$.
- is simple to implement, accurate when applied to Lane-Emden type equations and avoid tedious computational works.

The examples analyzed in the present paper reveal that the newly developed algorithms are easy, effective and accurate to solve the singular IVPs of Lane-Emden type equation.

30 2. Description of the method and its convergence

In this section, the VIM is described for solving Eq. (1). This method provides the solution as a sequence of iterations. It gives rapidly convergent successive approximations of the exact solution if such a solution exists, otherwise approximations can be used for numerical purposes.

The idea of the VIM is very simple and straightforward. To explain the basic idea of the VIM, we first consider Eq. (1) as follows:

$$L[y(x)] + N[y(x)] = g(x), \quad (2)$$

with

$$L[y(x)] = y''(x) + \frac{2}{x}y'(x) \quad \text{and} \quad N[y(x)] = f(x, y(x)), \quad (3)$$

where L denotes the linear operator with respect to y and N is a nonlinear operator with respect to y . The basic character of the VIM is to construct a correction functional according to the variational method as:

$$y_{n+1}(x) = y_n(x) + \int_0^x \lambda(t) \left(y_n''(t) + \frac{2}{t}y_n'(t) + f(t, \tilde{y}_n(t)) - g(t) \right) dt, \quad (4)$$

where λ is a general Lagrange multiplier, which can be identified optimally via variational theory, the subscript n denotes the n th approximation, and \tilde{y}_n is considered as a restricted variation, namely $\delta\tilde{y}_n = 0$. Successive approximations, $y_{n+1}(x)$, will be obtained by applying the obtained Lagrange multiplier and a properly chosen initial approximation $y_0(x)$. Consequently, the exact solution can be obtained by using

$$y(x) = \lim_{n \rightarrow \infty} y_n(x). \quad (5)$$

Now, to determine the optimal value of $\lambda(t)$, we continue as follows:

$$\delta y_{n+1}(x) = \delta y_n(x) + \delta \int_0^x \lambda(t) \left(y_n''(t) + \frac{2}{t}y_n'(t) \right) dt, \quad (6)$$

which the stationary conditions can be obtained from the relation (6) as:

$$\begin{cases} 1 - \lambda'(x) + \frac{2}{x}\lambda(x) = 0, \\ \lambda(x) = 0, \\ \lambda''(x) - 2\frac{x\lambda'(x) - \lambda(x)}{x^2} = 0, \end{cases} \quad (7)$$

and the Lagrange multiplier is gained as

$$\lambda(t) = -(t - \frac{t^2}{x}) \quad (8)$$

Finally, the iteration formula can be given as:

$$y_{n+1}(x) = y_n(x) - \int_0^x \left(t - \frac{t^2}{x}\right) \left(y_n''(t) + \frac{2}{t}y_n'(t) + f(t, y_n(t)) - g(t)\right) dt, \quad (9)$$

It is interesting to note that for linear Lane-Emden type equations, its exact solution can be obtained easily by only one iteration step due to the fact that the multiplier can be suitably identified, as will be shown in this paper later.

Now we will have the following proposition for the iteration formula (9).

Proposition 1. *If $y(x) \in C^2[0, T]$, then, for $x \leq T$*

$$\int_0^x \left(t - \frac{t^2}{x}\right) \left(y''(t) + \frac{2}{t}y'(t)\right) dt = y(x) - y(0). \quad (10)$$

PROOF. The left side of the relation (10) can be written as below:

$$\int_0^x \left(t - \frac{t^2}{x}\right) (y''(t)) dt + \int_0^x \left(2 - \frac{2t}{x}\right) (y'(t)) dt, \quad (11)$$

now integrating by parts first integral yields

$$\begin{aligned} & \left[t - \frac{t^2}{x}\right]_{t=0}^{t=x} - \int_0^x \left(1 - \frac{2t}{x}\right) (y'(t)) dt + \int_0^x \left(2 - \frac{2t}{x}\right) (y'(t)) dt \\ & = \int_0^x y'(t) dt = y(x) - y(0), \end{aligned} \quad (12)$$

this ends the proof of (10). \square

Thus, in the light of (9) and (10), therefore, we will have the following simple variational iteration formula:

$$y_{n+1}(x) = y(0) - \int_0^x \left(t - \frac{t^2}{x}\right) (f(t, y_n(t)) - g(t)) dt, \quad (13)$$

40 The VIM (13) makes a recurrence sequence $\{y_n(x)\}$. Obviously, the limit of this sequence will be the solution of (1) if this sequence is convergent.

In order to prove the sequence $\{y_n(x)\}$ is convergent, we construct a series

$$y_0(x) + [y_1(x) - y_0(x)] + \cdots + [y_n(x) - y_{n-1}(x)] + \cdots. \quad (14)$$

Noticing that

$$S_{n+1} = y_0(x) + [y_1(x) - y_0(x)] + \cdots + [y_n(x) - y_{n-1}(x)] = y_n(x), \quad (15)$$

the sequence $\{y_n(x)\}$ will be convergent if the series is convergent.

Theorem 1. *If $N[y(x)] = f(x, y)$ is Lipschitz-continuous in $[0, T]$ and $g(x) \in C[0, T]$ then the series of (14) is convergent, i.e., the sequence $\{y_n(x)\}$ is convergent for $x \in [0, T]$.*

PROOF. According to (13), note that

$$|y_1(x) - y_0(x)| = \left| \int_0^x \left(t - \frac{t^2}{x} \right) (f(t, y_0(t)) - g(t)) dt \right| \leq MNx, \quad (16)$$

where

$$M = \max_{0 \leq t \leq x \leq T} \left| t - \frac{t^2}{x} \right| \quad \text{and} \quad N = \max_{0 \leq t \leq x \leq T} |f(t, y_0(t)) - g(t)| \quad (17)$$

From (13) and (16), and the assumption that $|f(t, y_n) - f(t, y_{n-1})| \leq L|y_n - y_{n-1}|$ where L denotes the Lipschitz constant of $f(x, y)$, it follows that

$$|y_2(x) - y_1(x)| \leq ML \left| \int_0^x |y_1(t) - y_0(t)| dt \right| \leq \frac{N (MLx)^2}{L 2!}, \quad (18)$$

$$|y_3(x) - y_2(x)| \leq ML \left| \int_0^x |y_2(t) - y_1(t)| dt \right| \leq \frac{N (MLx)^3}{L 3!}, \quad (19)$$

⋮

$$|y_n(x) - y_{n-1}(x)| \leq \frac{N (MLx)^n}{L n!}. \quad (20)$$

In view of (20), the convergence of the series (14) can be concluded for the solution domain $x < T$. Therefore the series of (14) is absolute convergence, i.e., the sequence $\{y_n(x)\}$ is convergent for $x \in [0, T]$. \square

2.1. A truncated VIM

The successive iterations of the VIM may be very complex, so that the resulting integrals in the relation (4) may not be performed analytically. Also, the implementation of the VIM generally leads to calculation of unneeded terms, which more time is consumed in repeated calculations for series solutions. Here, an effective modification of the VIM is applied to eliminate these repeated calculations. To completely stop these repeats in each step, provided that the integrand of (4) in each of iterations is expanded in multivariate Taylor series around $x = 0$, we propose the following improvement of the VIM (4), which is called the truncated VIM (TV):

$$y_{n+1}(x) = y_n(x) - \int_0^x F_n(x, t) dt, \quad (21)$$

where

$$\left(t - \frac{t^2}{x}\right) \left(y_n''(t) + \frac{2}{t}y_n'(t) + f(t, y_n(t)) - g(t)\right) = F_n(x, t) + O(x^{n+1}) + O(t^{n+1}). \quad (22)$$

50 It is noteworthy to point out that the TV formula (21) can cancel all the repeated calculations and terms that are not needed as will be shown below. Furthermore, it can reduce the size of calculations. Most importantly, however, it is the fact that the TV algorithm (21) solves a Lane-Emden equation exactly if its solution is an algebraic polynomial up to some degree.

55 2.2. A local VIM

In general, by using the TV formula (21), we obtain a series solution, which in practice is a truncated series solution. This series solution gives a good approximation to the exact solution in a small region of x . An easy and reliable way of ensuring validity of the approximations (21) for large x is to determine the solution in a sequence of equal subintervals of x , i.e. $I_i = [x_i, x_{i+1}]$ where $h_i = x_{i+1} - x_i$, $i = 0, 1, \dots, N - 1$, with $x_0 = 0$ and $x_N = T$. According to the relation (21), therefore, we can construct the following piecewise TV

approximations (PTV) in the subintervals I_i . On $[x_0, x_1]$, let

$$\left\{ \begin{array}{l} y_{1,m+1}(x) = y_{1,m}(x) - \int_{x_0}^x F_{1,m}(x,t)dt, \quad m = 0, 1, \dots, n_1 - 1, \\ y_{1,0}(x) = y(0) + y'(0)(x - x_0) = c_0 + c'_0(x - x_0), \\ \left(t - \frac{t^2}{x} \right) \left(y''_{1,m}(t) + \frac{2}{t}y'_{1,m}(t) + f(t, y_{1,m}(t)) - g(t) \right) = F_{1,m}(x, t) \\ \quad + O((x - x_0)^{n+1}) + O((t - x_0)^{n+1}), \end{array} \right. \quad (23)$$

then we can obtain the n_1 -order approximation $y_{1,n_1}(x)$ on $[x_0, x_1]$. On $[x_1, x_2]$, let

$$\left\{ \begin{array}{l} y_{2,m+1}(x) = y_{2,m}(x) - \int_{x_1}^x F_{2,m}(x,t)dt, \quad m = 0, 1, \dots, n_2 - 1, \\ y_{2,0}(x) = y_{1,n_1}(x_1) + y'_{1,n_1}(x_1)(x - x_1) = c_1 + c'_1(x - x_1), \\ \left(t - \frac{t^2}{x} \right) \left(y''_{2,m}(t) + \frac{2}{t}y'_{2,m}(t) + f(t, y_{2,m}(t)) - g(t) \right) = F_{2,m}(x, t) \\ \quad + O((x - x_1)^{n+1}) + O((t - x_1)^{n+1}), \end{array} \right. \quad (24)$$

then we can obtain the n_2 -order approximation $y_{2,n_2}(x)$ on $[x_1, x_2]$. In a similar way, on $[x_i, x_{i+1}]$, $i = 2, 3, \dots, N - 1$, let

$$\left\{ \begin{array}{l} y_{i+1,m+1}(x) = y_{i+1,m}(x) - \int_{x_i}^x F_{i+1,m}(x,t)dt, \quad m = 0, 1, \dots, n_{i+1} - 1, \\ y_{i+1,0}(x) = y_{i,n_i}(x_i) + y'_{i,n_i}(x_i)(x - x_i) = c_i + c'_i(x - x_i), \\ \left(t - \frac{t^2}{x} \right) \left(y''_{i+1,m}(t) + \frac{2}{t}y'_{i+1,m}(t) + f(t, y_{i+1,m}(t)) - g(t) \right) = F_{i+1,m}(x, t) \\ \quad + O((x - x_i)^{n+1}) + O((t - x_i)^{n+1}), \end{array} \right. \quad (25)$$

then we can obtain the n_{i+1} -order approximation $y_{i+1,n_{i+1}}(x)$ on $[x_i, x_{i+1}]$.

Therefore, according to (23)-(25), the approximation of Eq. (1) on the entire interval $[0, T]$ can be obtained. It should be emphasized that the VIM and TV algorithms provide analytical solutions in $[0, T]$, while the PTV technique provides analytical solutions in $[x_i, x_{i+1}]$, which are continuous at the end points of each interval, i.e., $y_{i,n_i}(x_i) = c_i = y_{i+1,n_{i+1}}(x_i)$ and $y'_{i,n_i}(x_i) = c'_i = y'_{i+1,n_{i+1}}(x_i)$, $i = 1, 2, \dots, N - 1$.

It is obvious that the best PTV method of (25) can be achieved by using a variable order of n_{i+1} and a variable step size h_i in the solution to obtain

65 a specified tolerance. Therefore, the following adaptive strategy based on the variable step size is proposed for the PTV method, which we summarize it as the APTV (see, e.g., [6] and the references therein). This technique simplifies computation, and saves time and work, as will be observed later in this paper.

Let $\mathbf{y}_{i+1,k}$ be the solution of the fixed k -order PTV formula with the step size h_i and $\widehat{\mathbf{y}}_{i+1,k}$ the solution with the step size $h_i/2$. Taking the difference of $\mathbf{y}_{i+1,k}$ and $\widehat{\mathbf{y}}_{i+1,k}$, the local error estimator of $\mathbf{y}_{i+1,k}$

$$Est = \widehat{\mathbf{y}}_{i+1,k} - \mathbf{y}_{i+1,k}, \quad (26)$$

is defined. This value is an estimation of the main part of the local discretization error of the method. Additionally, let r be the dimension of the ODE system, and $Atol$ and $Rtol$ the user-specified absolute and relative error tolerances. The tolerances occurring in each step are denoted by

$$Tol_j = Atol + Rtol \cdot |y_{i+1,k}^j|, \quad j = 1, \dots, r. \quad (27)$$

Taking

$$err = \sqrt{\frac{1}{r} \sum_{j=1}^r \left(\frac{Est}{Tol_j} \right)^2}, \quad (28)$$

as a measure we find an optimal step size h_{opt} by comparing err to 1. Thus we obtain the optimal step size as

$$h_{opt} = h_i \cdot \left(\frac{1}{err} \right)^\alpha, \quad (29)$$

where for $err \leq fac_{err}$ ($fac_{err} \in (0, 1]$), we use $\alpha = \frac{1}{k+1}$, and for $err > fac_{err}$, $\alpha = \frac{1}{k}$. This is, of course, not the best choice for all problems. The new step size

$$h_{new} = h_{i+1} = h_i \cdot \min \left\{ fac_{max}, \max \left\{ fac_{min}, fac \cdot \left(\frac{1}{err} \right)^\alpha \right\} \right\}, \quad (30)$$

is obtained by using err with k as order of the approximation, instead of order
70 of consistency. The integration of the growth factors fac_{max} and fac_{min} to relation (30) prevents for too large step increase and contribute to the safety of the code. Additionally, using the safety factor fac makes sure that err will be

accepted in the next step with high probability. The step is accepted, in case that $err \leq fac_{err}$ otherwise it is rejected and then the procedure is redone. In
75 both cases the new solution is computed with h_{new} as step size.

3. Implementations

To give a clear overview of the content of this study, several Lane-Emden type equations will be studied. These equations will be tested by the above-mentioned algorithms, which will ultimately show the usefulness and accuracy
80 of these methods. Moreover, the numerical results indicate that the approach is easy to implement. All the results here are calculated by using the symbolic calculus software Maple 17. Also, all calculations are carried out in a Toshiba Tecra A8 (Windows 8.1 Professional): Intel(R) Core(TM)2 Duo Processor T7200 (2.00GHz, 4MB Cache, 997 MHz, 0.99 GB of RAM).

Example 1. As a first example, we consider the following linear, non-homogeneous Lane-Emden equation, i.e., Eq. (1) with $f(x, y) = y$ and $g(x) = 6 + 12x + x^2 + x^3$ (see, e.g., [12]):

$$y'' + \frac{2}{x}y' + y = 6 + 12x + x^2 + x^3, \quad (31)$$

subject to the initial conditions

$$y(0) = 0 \quad \text{and} \quad y'(0) = 0.$$

The VIM has a very simple approach. Its concepts begin with dividing the left hand (31) into two parts, i.e., the linear operator L and the nonlinear operator N as:

$$L[y(x)] = y'' + \frac{2}{x}y' + y \quad \text{and} \quad N[y(x)] \equiv 0. \quad (32)$$

This will allow us to construct a variational iteration relation for Eq. (31) as follows:

$$y_{n+1}(x) = y_n(x) - \int_0^x \left(\frac{t}{x} \sin(x-t) \right) \left(y_n''(t) + \frac{2}{t}y_n'(t) + y_n(t) - 6 - 12x - x^2 - x^3 \right) dt. \quad (33)$$

By using simple integration by parts, similar to Proposition 1, we will have

$$\int_0^x \left(\frac{t}{x} \sin(x-t) \right) \left(y_n''(t) + \frac{2}{t} y_n'(t) + y_n(t) \right) dt = y(x) - y(0) \frac{\sin(x)}{x}. \quad (34)$$

In the light of (33) and (34), therefore, we have the following VIM:

$$y_{n+1}(x) = y_0 \frac{\sin(x)}{x} + \int_0^x \left(\frac{t}{x} \sin(x-t) \right) (6 + 12x + x^2 + x^3) dt, \quad (35)$$

where $y_0 = y(0)$ and $y_0(x) = y(0) + y'(0)x$. According to (35), therefore, we get the following approximations with starting the initial guess $y_0(x) = 0$:

$$y_n(x) = x^2 + x^3 \quad \text{for all } n \geq 1, \quad (36)$$

85 which is the exact solution of the Lane-Emden equation (31). This proves our above-mentioned claim that the VIM could solve the linear Lane-Emden equation by only one iteration.

Example 2. As other example, we consider the nonlinear, non-homogeneous Lane-Emden equation, i.e., Eq. (1) with $f(x, y) = y^3$ and $g(x) = 6 + x^6$ (see, e.g., [12]):

$$y'' + \frac{2}{x} y' + y^3 = 6 + x^6, \quad (37)$$

subject to the initial conditions

$$y(0) = 0 \quad \text{and} \quad y'(0) = 0.$$

Here, we aim to solve the equation (37) by means of the TV algorithm (21). According to (21), we can easily obtain the following approximations of the TV with starting the initial approximation $y_0(x) = 0$:

$$\begin{aligned} y_1(x) &= 0, \\ y_n(x) &= x^2 \quad \text{for all } n \geq 2, \end{aligned} \quad (38)$$

which the TV algorithm yields the exact solution. This also demonstrates our above-noted claim that the PV algorithm could solve the linear/nonlinear Lane-
 90 Emden equation exactly if its solution is an algebraic polynomial up to some degree.

Example 3. As final example, we consider the nonlinear, homogeneous Lane-Emden-type equation, i.e., Eq. (1) with $f(x, y) = e^y$ and $g(x) = 0$ (see, e.g., [12]):

$$y'' + \frac{2}{x}y' + e^y = 0, \quad (39)$$

subject to the initial conditions

$$y(0) = 0 \quad \text{and} \quad y'(0) = 0.$$

Here, we aim to solve the equation (39) by means of the above-proposed methods. Since the integration of the nonlinear term e^y in Eq. (39) is not easily evaluated, thus the VIM requires a large amount of computational work to obtain few iterations of the solution (we can replace the nonlinear term with a series of finite components). However, we use the modified VIM method, i.e., the TV algorithm (21). According to (21), we can easily obtain the following approximations of (39) with starting the initial approximation $y_0(x) = 0$:

$$\begin{aligned} y_2(x) &= -\frac{1}{6}x^2, \\ y_4(x) &= -\frac{1}{6}x^2 + \frac{1}{120}x^4, \\ y_6(x) &= -\frac{1}{6}x^2 + \frac{1}{120}x^4 - \frac{1}{1890}x^6, \end{aligned} \quad (40)$$

and so on. Fig. 1 shows a comparison of approximation obtained using the 20th-order TV algorithm with the numerical solution of Eq. (39).

As observed, the TV algorithm (21) in solving Eq. (39) gives good approx-
 95 imations to the exact solution in a small region of x . In order to enlarge the convergence region of the series solution, here we implement the PTV (25) proposed in Section 2.2. According to (25), taking $N = 4000$ and $n_{i+1} = 4$, we can obtain the approximations of (36) on $[0, 1000]$. Fig. 2 shows the absolute error (the difference between the approximate value and the numerical value) of the
 100 PTV solution for $n_{i+1} = 4$ and $h_i = 0.25$. From Fig. 2, it is easily found that the present approximation is efficient for a larger interval.

Now, in order to show the efficiency of the above adaptive mechanism controlling the truncation error, we solve the above system using the before-mentioned APSP algorithm. The numerical results can be observed in Table 1. In Table 1,




Figure 1: Approximate solution for Example 3 using the TV algorithm where the dotted-line: the 20th-order TV algorithm and symbol: the numerical solution.




Figure 2: Shows the absolute error ($E(x) = |y_4(x) - y_{Numeric}(x)|$) of the 4th-order PTV solution for Example 3.

Table 1: The numerical results obtained from solving Example 3 using the 4th-order APTV algorithm when $fac_{err} = 1$, $fac = 0.9$, $fac_{min} = 0.5$ and $fac_{max} = 1.5$.

Algorithm	T	$Atol$	$Rtol$	No. of steps	CPU time (s)
APSP	1000	10^{-10}	10^{-10}	1030	4.156
APSP	1000	10^{-11}	10^{-11}	1819	7.047
APSP	1000	10^{-12}	10^{-12}	3223	12.156
APSP	1000	10^{-13}	10^{-13}	5720	21.765

105 we listed the costed number of steps (labeled as No. of steps) for some different values of T , $Atol$ and $Rtol$, and the corresponding costed CPU elapsed time (labeled as CPU time).

Moreover, in Fig. 3, one can see the plot of the variable step size using the fourth-order APTV algorithm for $Atol = Rtol = 10^{-13}$ under the assumptions 110 of Table 1. By observing this graph we can perfectly comprehend how the developed method works.

Furthermore, the local discretization error of the APTV algorithm for the value $Atol = Rtol = 10^{-13}$ under the assumptions of Table 1, which is an estimation of the principal portion of the local error, have been given in Fig. 4.

115 In closing our analysis, we point out that three concreted modeling equations of second-order singular IVPs of the Lane-Emden type equation were investigated by using the algorithms proposed in this paper, and the obtained results have shown noteworthy performance.

4. Conclusion

120 Application of the methods based on the VIM presented in this paper to three Lane-Emden type equations indicates that for linear Lane-Emden type equations, its exact solution, if such a solution exists, can be obtained easily by only one iteration step due to the fact that the multiplier can be suitably identified, that the TP algorithm can solve a nonlinear Lane-Emden differential 125 equation exactly if its solution is an algebraic polynomial up to some degree,




Figure 3: Variable step size of the 4th-order APTV algorithm when $Atol = Rtol = 10^{-13}$ for Example 3.




Figure 4: Local error of the 4th-order APTV algorithm when $Rtol = Atol = 10^{-13}$ for Example 3.

and that for nonlinear Lane-Emden type equations can be useful in general.

It is well-known that the achievement of methods to solve the nonlinear IVPs of ODEs depends on the use of adaptive step size mechanisms controlling the truncation error. For this reason, an adaptive version of the VIM was proposed.

130 The numerical results demonstrate that the VIM is a useful analytic tool for solving the Lane-Emden type equations.

References

- [1] Bender, C.M., Milton, K.A., Pinsky, S.S., Simmons L.M., 1989. A new perturbative approach to nonlinear problems, *J. Math. Phys.* 30, 1447–1455.
135
- [2] Chandrasekhar, S., 1967. *Introduction to the Study of Stellar Structure*. Dover, New York.
- [3] Chowdhury, M.SH., Hashim, I., 2007. Solutions of a class of singular second-order IVPs by homotopy–perturbation method, *Phys. Lett. A* 365, 439–447.
- 140 [4] Davis, H.T., 1962. *Introduction to Nonlinear Differential and Integral Equations*. Dover, New York.
- [5] Ghorbani, A., Momani, S., 2010. An effective variational iteration algorithm for solving Riccati differential equations. *Appl. Math. Lett.* 23, 922–927.
- 145 [6] Hairer, E., Wanner, G., 1991. *Solving Ordinary Differential Equations. II*. Springer-Verlag, Berlin.
- [7] He, J.H., 1999. Variational iteration method—a kind of non-linear analytical technique: some examples. *Int. J. Nonlinear Mech.* 34, 699–708.
- [8] He, J.H., 2003. Variational approach to the Lane-Emden equation. *Appl. Math. Comput.* 143, 539–541.
150

- [9] Krivec, R., Mandelzweig V.B., 2001. Numerical investigation of quasilinearization method in quantum mechanics. *Comput. Phys. Commun.* 138, 69–79.
- [10] Liao, SH.J., 2003. A new analytic algorithm of Lane-Emden type equations. *Appl. Math. Comput.* 142, 1–16.
- [11] Mandelzweig, V.B., Tabakin, F., 2001. Quasilinearization approach to nonlinear problems in physics with application to nonlinear ODEs. *Comput. Phys. Commun.* 141, 268–281.
- [12] Parand, K., Dehghan, M., Rezaei, A.R., Ghaderi, S.M., 2010. An approximation algorithm for the solution of the nonlinear LaneEmden type equations arising in astrophysics using Hermite functions collocation method. *Comput. Phys. Commun.* 181, 1096–1108.
- [13] Ramos, J.I., 2005. Linearization techniques for singular initial-value problems of ordinary differential equations. *Appl. Math. Comput.* 161, 525–542.
- [14] Ramos, J.I., 2008. Series approach to the Lane-Emden equation and comparison with the homotopy perturbation method. *Chaos Solitons Fractals* 38, 400–408.
- [15] Richardson, O.U., 1921. *The Emission of Electricity from Hot Bodies*. London.
- [16] Shawagfeh, N.T., 1993. Nonperturbative approximate solution for Lane-Emden equation. *J.Math. Phys.* 34, 4364–4369.
- [17] Wazwaz, A.M., 2001. A new algorithm for solving differential equations of Lane-Emden type. *Appl. Math. Comput.* 118, 287–310.
- [18] Yildirim, A., Ozis, T., 2007. Solutions of singular IVPs of Lane-Emden type by homotopy perturbation method. *Phys. Lett. A* 369, 70–76.